

Regulation of lipid accumulation in 3T3-L1 cells: insulin-independent and combined effects of fatty acids and insulin

T. A. Kokta¹, A. L. Strat¹, M. R. Papasani¹, J. I. Szasz¹, M. V. Dodson² and R. A. Hill^{1†}

¹Department of Animal and Veterinary Science, University of Idaho, PO Box 442330, Moscow, ID 83844-2330, USA; ²Department of Animal Sciences, Washington State University, Pullman, WA 99164, USA

(Received 12 January 2007; Accepted 1 August 2007)

The insulin-independent and combined effects of fatty acids (FA; linoleic and oleic acids) and insulin in modulating lipid accumulation and adipogenesis in 3T3-L1 cells was investigated using a novel protocol avoiding the effects of a complex hormone 'induction' mixture. 3T3-L1 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) plus serum (control) or in DMEM plus either 0.3 mmol/l linoleic or oleic acids with 0.3 mmol/l FA-free bovine serum albumin in the presence or absence of insulin. Cells were cultured for 4 to 8 days and cell number, lipid accumulation, peroxisome proliferator-activated receptor-gamma (PPAR- γ) and glucose transporter 4 (GLUT-4) protein expression were determined. Cell number appeared to be decreased in comparison with control cultures. In both oleic acid and linoleic acid-treated cells, notably in the absence (and presence) of insulin, oil-red O stain-positive cells showed abundant lipid. The percentage of cells showing lipid accumulation was greater in FA-treated cultures compared with control cells grown in DMEM plus serum ($P < 0.001$). Treatment with both linoleic and oleic acid-containing media evoked higher levels of PPAR- γ than observed in control cultures ($P < 0.05$). GLUT-4 protein also increased in response to treatment with both linoleic and oleic acid-containing media ($P < 0.001$). Lipid accumulation in 3T3-L1 cells occurs in response to either oleic or linoleic acids independently of the presence of insulin. Both PPAR- γ and GLUT-4 protein expression were stimulated. Both proteins are considered markers of adipogenesis, and these observations suggest that these cells had entered the physiological state broadly accepted as differentiated. Furthermore, 3T3-L1 cells can be induced to accumulate lipid in a serum-free medium supplemented with FA, without the use of induction protocols using complex hormone mixtures. We have demonstrated a novel model for the study of lipid accumulation that will improve the understanding of adipogenesis in adipocyte lineage cells.

Keywords: adipogenesis, fatty acids, GLUT-4, insulin, PPAR- γ

Introduction

Progression to lipid storage in adipocytes is characterised by two distinct phases. In the first step of hyperplastic expansion, progenitor cells are thought to become committed to the adipocyte lineage, after which they cannot revert back to a less differentiated 'stem-like' cell (Thompson *et al.*, 1998; Boone *et al.*, 2000). Once committed, adipoblasts undergo an exponential replication phase that terminates and the cell cycle arrests at gap 1 (G1). Early markers of differentiation, such as lipoprotein lipase (LPL) are then expressed, and these cells, known as preadipocytes, may then undergo proliferation (Boone *et al.*, 2000). After preadipocytes stop proliferating, late markers of differentiation, such as glycerol-3-phosphate

dehydrogenase (GPDH) and fatty acid synthetase (FAS) are detected. Cells then begin lipid accumulation in the cytosol at which time cells are termed adipocytes (Boone *et al.*, 2000).

Much of our understanding of these processes has developed through the use of *in vitro* models. A common approach when using *in vitro* models is to culture cells thought (or previously shown) to express adipogenesis lineage genes until they become confluent and then to apply a hormonal or xenobiotic agent 'cocktail' to 'induce' adipogenesis. Typically, these induction protocols include moderate to high concentrations of factors such as dexamethasone, methylisobutylxanthine and insulin in serum-containing media (Wu *et al.*, 1996; Sakaue *et al.*, 1998; Habinowski and Witters, 2001; Croissandeau *et al.*, 2002; Floyd and Stephens, 2003; Shin *et al.*, 2003). Both oleic (Guo *et al.*, 2000) and linoleic (Kang *et al.*, 2003) acids

† E-mail: rodhill@uidaho.edu

were reported to induce differentiation in 3T3-L1 preadipocytes. However, octanoate appeared to attenuate differentiation in 3T3-L1 cells (Han *et al.*, 2002). Conjugated linoleic acids have also been reported to inhibit differentiation in 3T3-L1 cells (Brodie *et al.*, 1999; Kang *et al.*, 2003).

One potential mechanism of enhanced adipogenesis from fatty acids (FA) may potentially be via peroxisome proliferator-activated receptors (PPARs) (Tontonoz *et al.*, 1994a and 1994b; Grimaldi, 2001; Bishop-Bailey and Wray, 2003). During differentiation of preadipocytes to adipocytes, PPAR- γ , PPAR- α and CCAAT/enhancer-binding protein factors (C/EBP) α , β , δ and ζ are important transcription factors involved in the regulation of this mechanism (Lee *et al.*, 1999; Kersten *et al.*, 2000; Lacasa *et al.*, 2001); PPAR- γ being activated prior to the activation of C/EBP- α . PPAR- γ is also required for survival of mature adipocytes (Imai *et al.*, 2004), and is activated by a variety of factors, including glucocorticoids (Wu *et al.*, 1996), unsaturated FA such as oleic acid, linoleic acid, eicosapentaenoic acid and arachidonic acid (Amri *et al.*, 1995) and insulin-sensitising reagents such as thiazolidinidiones (Spiegelman, 1998; Torii *et al.*, 1998; Yamauchi *et al.*, 2001; Li and Lazar, 2002). Hence, insulin is thought to be a factor in the increased expression of PPAR- γ . PPAR- γ activation also results in the initiation of expression of GLUT-4. This process also requires additional unknown factors, and appears to complete the regulatory pathway controlling glucose uptake in response to insulin (Wu *et al.*, 1998).

Normal fasting concentrations of total plasma free FA fall approximately in the range 0.3 to 0.6 mmol/l and during exercise may be as high as 1.5 mmol/l. The contributions of each FA to total fasting FA is variable, but the major contributors are palmitic, 16:0 (20 to 25 mass %), oleic, 18:1 (15 to 25 mass %) and linoleic acids, 18:2 (22 to 30 mass %). *In vivo*, peripheral tissues, including adipose will utilise free FA from plasma and liberated FA from triglycerides facilitated by vascular endothelium LPL activity; the uptake of FA by adipocytes being favoured by the local concentration gradient. Thus, during periods of high nutrition (high energy intake) or exercise, peripheral tissues will be exposed to FA concentrations equal to or higher than the range shown above. It is difficult to emulate this model *in vitro*, but to better understand the mechanisms of utilisation of FA in cellular lipid accumulation, it may be necessary to provide FA in this physiological concentration range when performing *in vitro* studies.

In the present study, we report induction of lipid accumulation in 3T3-L1 cells. We asked the question: can 3T3-L1 cells be induced to accumulate lipid and ultimately progress through the process of adipogenesis in the absence of the effects of a hormone 'cocktail'? The underlying hypothesis to the study was that supply of FA at physiological or slightly hyper-physiological concentrations would induce lipid accumulation and adipogenesis in confluent 3T3-L1 cells. This approach also facilitated investigation of the interaction of insulin effects. Thus, we also evaluated the

response of this model to the presence or absence of insulin. To provide a model we considered physiologically appropriate in terms of energy substrate and lipid precursors, a standard defined medium (Dulbecco's modified Eagle's medium, DMEM), supplemented with individual FA at a concentration approximating total FA concentration in serum (0.3 mmol/l), protected by the addition of FA-free bovine serum albumin (BSA) (0.3 mmol/l) was used. We report the effects of treatment on cell number, cellular lipid accumulation and expression of adipogenic markers PPAR- γ and GLUT-4 proteins in this model.

Material and methods

Material

Cell-culture plasticware was supplied by Nunc (Naperville, IL, USA). DMEM, foetal bovine serum (FBS), penicillin-streptomycin, gentamicin, insulin and trypsin were purchased from Gibco BRL (Rockville, MD, USA). BSA (FA free), insulin, oil-red O stain, oleic and linoleic acid adsorbed to BSA were obtained from Sigma (St Louis, MO, USA). Rabbit anti-PPAR- γ (RDi-PPARGabrx) was obtained from Research Diagnostics, Inc. (Flanders, NJ, USA). Rabbit anti-GLUT-4 antibody (GLU-T4ab654) was obtained from Abcam (Cambridge, MA, USA). Secondary antibody was Alexa Fluor 680 goat anti-rabbit IgG obtained from Molecular Probes (Eugene, OR, USA).

Cell culture

3T3-L1 preadipocytes were plated into 24-well plates at a density of 80 000 cells per well, and cultured at 37°C, 5% CO₂ in DMEM + 10% FBS + antibiotics. The confluent cultures (2 to 3 days) were then rinsed three times with DMEM and transferred to a defined medium treatment, or control cultures were maintained in DMEM + 10% FBS + antibiotics. Media were changed every 48 h. Parallel cultures were grown in T-75 flasks for protein studies. Cell lysates were obtained at days 4 and 8 of each treatment.

Cells were cultured in DMEM with either 0.3 mmol/l oleic or linoleic acid plus 0.3 mmol/l FA-free BSA, and either in the presence or absence of 200 nmol/l insulin. FAs were purchased (non-covalently) adsorbed to BSA, protecting cells from the effects of unbound FA and providing similar FA availability to that *in vivo*.

Cell number and cell morphology

Cell number was determined by counting cells in each of 10 fields. Cultures from 24-well plates were fixed with 10% formalin and stained with oil-red O, to detect the presence of lipid, and counterstained with Harris haematoxylin.

Preparation of cell lysates and Western blot

Cells were freeze-fractured in liquid nitrogen, and cell lysates prepared in a buffer consisting of 50 mmol/l Tris, 1% NP-40, 0.25% sodium deoxycholate, 150 mmol/l NaCl, 1 mmol/l ethylenediamine tetraacetic acid (EDTA), 1 mmol/l

sodium vanadate, plus Complete Mini-Cocktail (Roche Diagnostics; GmbH, Germany) pH 7.3 (Hill *et al.*, 2004). Total protein was determined using the BCA assay (Pierce Chemical Co., Rockford, IL, USA) and 20 μ g protein was loaded per lane on SDS PAGE. Resolved proteins were transferred to nitrocellulose and detected in Western blot using (1/1000) polyclonal rabbit anti-PPAR- γ antibody or rabbit anti-GLUT-4 antibody, and secondary antibody (1/5000) Alexa Fluor 680 goat anti-rabbit IgG and scanned on an infra-red fluorescence imager (Odyssey; LI-COR, Inc., Lincoln, NE, USA).

Design and statistical analyses

Studies were designed as reported by others using cell line models (Chiaromonte *et al.*, 1998; Schneider *et al.*, 2001). Thus, each experiment was replicated using triplicate (or a greater number of replicate) assays from duplicate cultures for each test condition. Statistical analyses were conducted using the Statistical Analysis Systems Institute (2002). Lipid accumulation data were treated as binomial responses. Accordingly, cell counts were logit transformed, and analysed using a binomial probability distribution in PROC GENMOD. The model included effects of day (4 and 8), FA, insulin and the resulting interactions. Total cell numbers, PPAR- γ and GLUT-4 protein expression were analysed using the GLM procedure with a model including effects of day, treatment, and a day-by-treatment interaction. A single degree of freedom contrast was tested to compare controls with cultures treated with FA and insulin. Additional single degree of freedom contrasts were used to test the main effects of FA and insulin, and the resulting interaction. When a FA \times insulin \times day interaction was observed, main effects of FA and insulin and the resulting interaction were tested by day.

Results

Cell number

At day 4 of incubation, cell number per field of view was lower ($P < 0.05$) for insulin – compared with non-insulin-treated cultures and for oleic acid – compared with linoleic acid-treated cultures (Figure 1). A FA \times insulin interaction was noted ($P < 0.05$) on day 8 of incubation in which there was a greater number of cells in cultures treated with oleic acid plus insulin compared with cultures treated with oleic acid alone or with linoleic acid plus insulin. It should be noted that one factor contributing to change in cell number was mean cell size. Although cell size was not quantified, it may be assumed that cells filling with lipid become larger and thus cell number expressed as cells/field will decrease.

Lipid accumulation

Qualitatively, lipid accumulation was difficult to detect in control cells cultured in DMEM + 10% FBS (Figure 2). In both oleic acid- and linoleic acid-treated cells, in the presence and absence of insulin on days 4 and 8, oil-red O stain-positive cells showed abundant lipid.




Figure 1 3T3-L1 cells were cultured in 0.3 mmol/l oleic or linoleic acid, with or without 200 nmol/l insulin, for up to 8 days. Control cells were cultured in Dulbecco's modified Eagle's medium + 10% foetal bovine serum. For each treatment, cells were cultured in duplicate and 10 fields per culture were enumerated. Data show cell number per field (s.e.) for each treatment group cultured for (a) 4 days and (b) 8 days. There was a significant fatty acid (FA) effect on day 4 (a, b, c, $P < 0.05$) and a FA \times insulin interaction on day 8 ($P < 0.05$). Also on day 4, there was a significant effect of the presence or absence of insulin (m, n and r, s, each $P < 0.05$).

Quantitatively, the percentage of cells showing lipid accumulation was greater in FA-treated cultures compared with control cells grown in DMEM plus serum ($P < 0.001$). At day 4 of incubation, the percentage of cells showing lipid when cultured in FA in the absence of insulin, was lower compared with those cultured in FA in the presence of insulin ($P < 0.01$, Figure 3a). Also on day 4, the percentage of cells showing lipid, when cultured in linoleic acid was lower compared with those cultured in oleic acid ($P < 0.01$). On day 8, a FA \times insulin interaction was noted ($P < 0.05$) in which lipid accumulation was greatest ($P < 0.05$) for cultures containing oleic acid plus insulin (Figure 3b). Additionally, cultures containing oleic acid alone showed greater ($P < 0.05$) lipid accumulation compared with cultures containing linoleic acid with or without insulin (Figure 3b).

PPAR- γ protein expression

Treatment with both linoleic and oleic acid-containing media evoked higher levels of PPAR- γ than observed in DMEM plus serum on both days 4 and 8 ($P < 0.05$, Figure 4a and b). There were no differential effects of FA or insulin on




Figure 2 3T3-L1 cells were cultured, either in control medium (Dulbecco's modified Eagle's medium (DMEM) + 10% foetal calf serum) or in DMEM plus bovine serum albumin adsorbed fatty acids (0.3 mmol/l) as follows: either, linoleic acid, linoleic acid plus 200 nmol/l insulin, oleic acid or oleic acid plus 200 nmol/l insulin, and fixed and stained with oil-red O, after either 4 or 8 days.

PPAR- γ on day 4 (Figure 4a). After 8 days of culture, each FA showed a similar pattern in which the presence of insulin tended to increase PPAR- γ ($P = 0.1$, Figure 4b).

GLUT-4 protein expression

GLUT-4 protein appeared to be close to background levels in Western blots of cell lysates from control (DMEM plus serum) cultures. Treatment with both linoleic and oleic acid-containing media evoked higher levels of GLUT-4 than observed in DMEM plus serum on both days 4 and 8 ($P < 0.001$, Figure 5a and b). There was no effect of insulin on GLUT-4 on day 4, reflecting a similar observation of PPAR- γ . However, there was a FA effect in which cells cultured in linoleic acid showed higher GLUT-4 protein expression than cells cultured in oleic acid ($P < 0.05$, Figure 5a). Similar results were observed after 8 days of culture, in which there was no effect of insulin, and cells cultured in linoleic acid tended to show higher GLUT-4 expression than those cultured in oleic acid ($P = 0.1$, Figure 5b).

Discussion

Cell number was lower in cultures treated with FA alone and FA plus insulin in comparison to the controls.




Figure 3 3T3-L1 cells were cultured in 0.3 mmol/l oleic or linoleic acid, with or without 200 nmol/l insulin, for up to 8 days. Control cells were cultured in Dulbecco's modified Eagle's medium (DMEM) + 10% foetal bovine serum. For each treatment, cells were cultured in duplicate and 10 fields per culture were enumerated. Data are the percentage of cells showing lipid accumulation for each treatment group. (a) Day 4, the percentage of cells showing lipid accumulation was greater in fatty acid (FA)-treated cultures compared with control cells grown in DMEM plus serum ($P < 0.001$). Main effects of FA and insulin: the percentage of cells showing lipid, when cultured in FA in the absence of insulin, was lower compared with those cultured in FA in the presence of insulin ($P < 0.01$), and when cultured in linoleic acid was lower compared with those cultured in oleic acid ($P < 0.01$). Bars with different superscripts differ, oleic v. linoleic acid (insulin effect not denoted). (b) Day 8, the percentage of cells showing lipid accumulation was greater in FA-treated cultures compared with control cells grown in DMEM plus serum ($P < 0.001$). There was a FA \times insulin interaction: lipid accumulation was greatest ($P < 0.05$) for cultures containing oleic acid plus insulin. Cultures containing oleic acid alone showed greater ($P < 0.05$) lipid accumulation compared with cultures containing linoleic acid with or without insulin.

Presumably, FA treatment results in withdrawal from cell cycle, whereas cells in control media continued to proliferate. These data are consistent with a study of human preadipocytes by McNeel *et al.* (2003) who observed that the addition of either oleic acid or conjugated linoleic acid (in the presence of foetal calf serum (FCS), insulin, hydrocortisone and transferrin) resulted in cell number being 30% lower than in controls.

In the present study, effects of FA on cell number were not fully elucidated, and different FA may potentially have different effects. Others have found that oleic acid was better tolerated than palmitic acid, and may even have a




Figure 4 3T3-L1 cells were cultured in 0.3 mmol/l oleic or linoleic acid, with or without 200 nmol/l insulin, for up to 8 days. Control cells were cultured in Dulbecco's modified Eagle's medium+10% foetal bovine serum. For each treatment, cells were cultured in duplicate and peroxisome proliferator-activated receptor-gamma (PPAR- γ) protein was quantified (triplicate assays per culture) in Western blots of cell lysates using scanning laser excitation fluorescence (Odyssey, LI-COR). Data are expressed as pixels per mm² (s.e.), from 20 μ g protein per lane. (a) Day 4, overall treatment effect, PPAR- γ was increased v. control, ($P < 0.001$). There was no insulin or fatty acid (FA) effect. (b) Day 8, overall treatment effect, PPAR- γ was increased v. control ($P < 0.001$). Each FA showed a similar pattern in which the presence of insulin tended to increase PPAR- γ ($P = 0.1$).

role in rescuing cells from apoptosis in mouse pancreatic β -cells (Listenberger *et al.*, 2003) and rat ventricular myocytes (De Vries *et al.*, 1997) through the promotion of triglyceride accumulation instead of apoptotic pathways or alterations of the lipid membrane as seen by saturated FA. Evidence of apoptosis was not detected in the present study, caspase-3 activity being low and invariable across all treatments (data not shown). Another aspect of enumeration of these cells is the inevitable increase in cell size with lipid accumulation. Thus, part of the apparent decrease in cell number (expressed here as cells/field) may be attributed to larger cells occupying each field, and not loss of cells through mechanisms of cell death.

As most of the protocols used to study lipid accumulation of preadipocytes use high concentrations of insulin, there appears to be a lack of documented studies on the effects of the presence or absence of insulin in conjunction with physiological concentrations of FA. A novel finding of the present study is that treatment with oleic or linoleic acid, independent of insulin, was sufficient to induce lipid




Figure 5 3T3-L1 cells were cultured in 0.3 mmol/l oleic or linoleic acid, with or without 200 nmol/l insulin, for up to 8 days. Control cells were cultured in Dulbecco's modified Eagle's medium+10% foetal bovine serum. For each treatment, cells were cultured in duplicate and GLUT-4 protein was quantified (triplicate assays per culture) in Western blots of cell lysates using scanning laser excitation fluorescence (Odyssey, LI-COR). Data are expressed as pixels per mm² (s.e.) from 20 μ g protein per lane. (a) Day 4, overall treatment effect, GLUT-4 was increased v. control, ($P < 0.001$). There was no insulin effect. Protein expression of GLUT-4 was higher in cells cultured in linoleic acid compared with oleic acid-containing media ($P < 0.05$). (b) Day 8, data were similar to day 4, for the overall treatment effect, GLUT-4 was increased v. control ($P < 0.001$). There was no insulin effect, and cells cultured in linoleic acid tended to show higher GLUT-4 expression than those cultured in oleic acid-containing media ($P = 0.1$).

accumulation in 3T3-L1 cells. Many studies of adipocyte lineage cells report the addition of various hormone cocktails either with FCS (Brodie *et al.*, 1999; Guo *et al.*, 2000; Kang *et al.*, 2003; McNeel and Mersmann, 2003; McNeel *et al.*, 2003) or without FCS (Ding and Mersmann, 2001; Ding *et al.*, 2002; Hsu and Ding, 2003; Hutley *et al.*, 2003) in addition to FA to promote differentiation. These studies report that oleic and linoleic acids promote lipid accumulation in adipocytes. Oleic acid increased differentiation over controls in 3T3-L1 cells (Guo *et al.*, 2000), and human preadipocytes (McNeel *et al.*, 2003). Linoleic acid has been reported to increase differentiation over control in human preadipocytes (Brown and McIntosh, 2003; Hutley *et al.*, 2003) and 3T3-L1 adipocytes (Kang and Pariza, 2001; Kang *et al.*, 2003). In each of these studies, cells were induced to differentiate with a hormone 'cocktail', and the resulting cultures were continuously maintained in media containing either no serum, but including a hormone 'cocktail' (Hutley *et al.*, 2003), or FCS plus or minus FA and apparently high

concentrations of insulin for a further 2 to 12 days (Kang and Pariza, 2001; Hutley *et al.*, 2003; Kang *et al.*, 2003).

In the present study, oleic acid appeared to be more effective than linoleic acid in inducing lipid accumulation, with over 90% of cells accumulating lipid in response to oleic acid in comparison to about 60% of cells in response to linoleic acid. Oleic acid treatment retained higher efficacy on both days 4 and 8. This is apparently a more rapid response, although somewhat consistent with reports in the literature where oleic acid (plus FCS and hormone 'cocktail')-treated human adipocytes attained 90% differentiation by day 17 (McNeel *et al.*, 2003), while another study reported about 65% differentiation in human preadipocytes treated with linoleic acid (plus hormone 'cocktail') on day 21 (Hutley *et al.*, 2003). It should be noted that each of these studies used primary cultures, and the remaining undifferentiated cells may have been non-adipocyte lineage cell types. In the present study, there were no differences in lipid accumulation between linoleic acid alone and linoleic acid plus insulin-treated cultures on either day 4 or day 8. Thus, perhaps surprisingly, it appears that insulin is not required to induce or maintain lipid accumulation in this model. The percentage of cells containing lipid in response to FA observed on day 4 of culture was only slightly modified by day 8, although the size and number of lipid droplets in each treatment may have increased (Figure 2, not quantified). Although insulin has been widely used in other studies of adipocyte development, in the present study, its addition to each of the FA media, had only a numerically small effect on lipid accumulation. Analysed as an overall main effect, the addition of insulin was significant (Figure 3). However, the numerically small effect observed suggests that in this *in vitro* model, insulin may be much less important for induction of lipid accumulation than indicated by other *in vitro* models.

PPAR- γ protein expression was increased by FA or FA plus insulin treatment by 2.5- to 4.5-fold. These data are consistent with a study by Kang *et al.* (2003) in 3T3-L1 cells where these authors reported that PPAR- γ expression was increased on days 4 and 7 compared with day 2 in response to linoleic acid. However, the additional effect of the use of a hormone 'cocktail' in the medium prevents a direct comparison with the data of the present study.

Further evidence that 3T3-L1 cells cultured in the presence of FA had entered the physiological state considered to be adipogenesis was demonstrated by greatly increased expression of GLUT-4. This observation is consistent with other studies in 3T3-L1 cells (El-Jack *et al.*, 1999; Gross *et al.*, 2004) in transition from preadipocyte to adipocyte morphology. Interestingly, in the present study no insulin effect on total cellular GLUT-4 expression was observed as might have been expected; although it may be assumed that GLUT-4 would be recruited to the plasma membrane in response to insulin. In all previously reported studies of GLUT-4 expression in adipocytes, insulin was used for the induction of lipid accumulation. Hence, the present study is the first to report the independent effects of FA (or FA plus

insulin) on GLUT-4 in the absence of prior exposure to insulin. It appears that under the present experimental conditions, between culture days 4 and 8, these cells may be transitioning from preadipocytes to adipocytes. Thus, PPAR- γ and GLUT-4 are both expressed but additional, unknown intracellular regulatory mechanisms may not be fully activated.

Additionally, the effects of different FA on GLUT-4 expression have not yet been reported. We found linoleic acid to be initially more effective than oleic acid in inducing GLUT-4 expression. The differential response to linoleic and oleic acids was opposite in terms of lipid accumulation and GLUT-4 expression. This observation suggests that while oleic acid may be more lipogenic, linoleic acid up-regulated glucose transport, possibly reflecting a higher demand for ATP from glycolysis in its incorporation into lipid. The GLUT-4 response to oleic and linoleic acids is intriguing and suggests a direct PPAR- γ mediated mechanism. Further study is needed to clarify this issue. These data may be surprising, but indicate that insulin may not be essential for induction of lipid accumulation, or increased protein levels of PPAR- γ or GLUT-4.

This appears to be the first study to report lipid accumulation in adipocyte lineage cells in response to physiological concentrations of FA, in the absence of the influence of a complex hormone 'cocktail'. Linoleic and oleic acids are considered to be ligands for PPAR- γ (Amri *et al.*, 1995; Forman *et al.*, 1997; Margetic *et al.*, 2002; Ferré, 2004; Kokta *et al.*, 2004), however there appear to be no other *in vitro* adipocyte lineage cell studies, which have investigated PPAR- γ expression or lipid accumulation in the absence of insulin. The strategy of the current study was to provide FA in the media at approximately physiological concentrations, as building blocks for lipid synthesis. Because FA form a regular component of the extracellular milieu *in vivo*, the interpretation of regulation of PPAR- γ expression in the absence of FA seems to be biologically irrelevant. In the present study, the effect of physiological concentrations of FA in the culture media on FA concentration in the intracellular or nuclear compartments were not investigated. This information would clarify the potential roles of these FA as PPAR- γ ligands.

Using the conventional hormone 'cocktail' adipogenesis induction approach, two pathways are rapidly activated in preadipocytes: in response to elevated cyclic AMP levels stimulated by methylisobutyl xanthine or glucocorticoids activating glucocorticoid response elements, expression of C/EBP- β and C/EBP- δ is rapidly increased while insulin signals to activate PPAR- γ transcription, via the 3-phosphoinositide-dependent protein kinase-1 (PKD-1) pathway following treatment (Ferré, 2004). An additional pathway regulating early events in differentiation appears to respond rapidly to FA and is regulated via PPAR- δ (Forman *et al.*, 1997). This pathway is independent of insulin signalling to activate PPAR- γ . It also appears that these pathways may be at least partially redundant (Grimaldi, 2001), and implies that insulin may not be essential for adipogenesis. The

present study suggests that FA alone may induce the PPAR- δ pathway leading to progression of both PPAR- γ and GLUT-4 expression. Investigation of this hypothesis would require comprehensive study of role of FA in both the PPAR- δ and insulin-mediated pathways.

Conclusion

It appears that lipid accumulation occurs in response to either oleic or linoleic acids, and that both may act to increase PPAR- γ and GLUT-4 protein expression. These effects are independent of insulin. However, insulin appears to have a small enhancing effect on both lipid accumulation and PPAR- γ protein expression. Furthermore, 3T3-L1 cells can be induced to accumulate lipid in parallel to increased PPAR- γ and GLUT-4 protein expression in a serum-free medium supplemented with FA, without the use of typical induction protocols. The present report provides the basis of a novel model for the study of lipid accumulation and regulation of adipogenesis in adipocyte lineage cells. Because a complex hormone mix is not required, and energy and molecular building blocks are provided near to physiological levels for lipid synthesis, this model will provide additional insight into the regulation of adipogenesis.

Acknowledgements

This study was supported in part by National Research Initiative Competitive Grant numbers 2003-35206-12880 and 2003-35206-13617 from the USDA Cooperative State Research, Education and Extension Service.

References

Amri EZ, Bonino F, Ailhaud G, Abumrad NA and Grimaldi PA 1995. Cloning of a protein that mediates transcriptional effects of fatty acids in preadipocytes. Homology to peroxisome proliferator-activated receptors. *Journal of Biological Chemistry* 270, 2367–2371.

Bishop-Bailey D and Wray J 2003. Peroxisome proliferator-activated receptors: a critical review on endogenous pathways for ligand generation. *Prostaglandins and Other Lipid Mediators* 71, 1–22.

Boone C, Mourot J, Gregoire F and Remacle C 2000. The adipose conversion process: regulation by extracellular and intracellular factors. *Reproduction, Nutrition, Development* 40, 325–358.

Brodie AE, Manning VA, Ferguson KR, Jewell DE and Hu CY 1999. Conjugated linoleic acid inhibits differentiation of pre- and post-confluent 3T3-L1 preadipocytes but inhibits cell proliferation only in pre-confluent cells. *Journal of Nutrition* 129, 602–606.

Brown JM and McIntosh MK 2003. Conjugated linoleic acid in humans: regulation of adiposity and insulin sensitivity. *Journal of Nutrition* 133, 3041–3046.

Chiaromonte R, Bartolini E, Testolin C and Comi P 1998. Regulation of the human GLUT4 gene expression in tumor RD18 cell line. *Pathobiology* 66, 191–195.

Croissant-deau G, Chretien M and Mbikay M 2002. Involvement of matrix metalloproteinases in the adipose conversion of 3T3-L1 preadipocytes. *Biochemical Journal* 364, 739–746.

De Vries J, Vork M, Roemen T, De Jong Y, Cleutjens J, Van der Vusse G and Van Bilsen M 1997. Saturated but not mono-unsaturated fatty acids induce apoptotic cell death in neonatal rat ventricular myocytes. *Journal of Lipid Research* 38, 1384–1394.

Ding ST and Mersmann HJ 2001. Fatty acids modulate porcine adipocyte differentiation and transcripts for transcription factors and adipocyte-characteristic proteins. *Journal of Nutritional Biochemistry* 12, 101–108.

Ding ST, McNeel RL and Mersmann HJ 2002. Modulation of adipocyte determination and differentiation-dependent factor 1 by selected polyunsaturated fatty acids. *In vitro Cellular and Developmental Biology – Animal* 38, 352–357.

El-Jack AK, Kandror KV and Pilch PF 1999. The formation of an insulin-responsive vesicular cargo compartment is an early event in 3T3-L1 adipocyte differentiation. *Molecular Biology of the Cell* 10, 1581–1594.

Ferré P 2004. The biology of peroxisome proliferator-activated receptors: relationship with lipid metabolism and insulin sensitivity. *Diabetes* 53 (Suppl. 1), S43–S50.

Floyd ZE and Stephens JM 2003. STAT5A promotes adipogenesis in nonprecursor cells and associates with the glucocorticoid receptor during adipocyte differentiation. *Diabetes* 52, 308–314.

Forman BM, Chen J and Evans RM 1997. Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proceedings of the National Academy of Sciences of the United States of America* 94, 4312–4317.

Grimaldi PA 2001. The roles of PPARs in adipocyte differentiation. *Progress in Lipid Research* 40, 269–281.

Gross DN, Farmer SR and Pilch PF 2004. Glut4 storage vesicles without Glut4: transcriptional regulation of insulin-dependent vesicular traffic. *Molecular and Cellular Biology* 24, 7151–7162.

Guo W, Choi JK, Kirkland JL, Corkey BE and Hamilton JA 2000. Esterification of free fatty acids in adipocytes: a comparison between octanoate and oleate. *Biochemical Journal* 349, 463–471.

Habinowski SA and Witters LA 2001. The effects of AICAR on adipocyte differentiation of 3T3-L1 cells. *Biochemical and Biophysical Research Communications* 286, 852–856.

Han J, Farmer SR, Kirkland JL, Corkey BE, Yoon R, Pirtskhalava T, Ido Y and Guo W 2002. Octanoate attenuates adipogenesis in 3T3-L1 preadipocytes. *Journal of Nutrition* 132, 904–910.

Hill RA, Strat AL, Hughes NJ, Kokta TJ, Dodson MV and Gertler A 2004. Early insulin signaling cascade in a model of oxidative skeletal muscle: mouse Sol8 cell line. *Biochimica et Biophysica Acta* 1693, 205–211.

Hsu JM and Ding ST 2003. Effect of polyunsaturated fatty acids on the expression of transcription factor adipocyte determination and differentiation-dependent factor 1 and of lipogenic and fatty acid oxidation enzymes in porcine differentiating adipocytes. *British Journal of Nutrition* 90, 507–513.

Hutley LJ, Newell FM, Joyner JM, Suchting SJ, Herington AC, Cameron DP and Prins JB 2003. Effects of rosiglitazone and linoleic acid on human preadipocyte differentiation. *European Journal of Clinical Investigation* 33, 574–581.

Imai T, Takakuwa R, Marchand S, Dentz E, Bornert JM, Messaddeq N, Wendling O, Mark M, Desvergne B, Wahli W, Chambon P and Metzger D 2004. Peroxisome proliferator-activated receptor gamma is required in mature white and brown adipocytes for their survival in the mouse. *Proceedings of the National Academy of Sciences of the United States of America* 101, 4543–4547.

Kang K and Pariza MW 2001. *trans*-10, *cis*-12-conjugated linoleic acid reduces leptin secretion from 3T3-L1 adipocytes. *Biochemical and Biophysical Research Communications* 287, 377–382.

Kang K, Liu W, Albright KJ, Park Y and Pariza MW 2003. *trans*-10, *cis*-12 CLA inhibits differentiation of 3T3-L1 adipocytes and decreases PPAR gamma expression. *Biochemical and Biophysical Research Communications* 303, 795–799.

Kersten S, Mandard S, Tan NS, Escher P, Metzger D, Chambon P, Gonzalez FJ, Desvergne B and Wahli W 2000. Characterization of the fasting-induced adipose factor FIAF, a novel peroxisome proliferator-activated receptor target gene. *Journal of Biological Chemistry* 275, 28488–28493.

Kokta TA, Dodson MV, Gertler A and Hill RA 2004. Intercellular signaling between adipose tissue and muscle tissue. *Domestic Animal Endocrinology* 27, 303–331.

Lacasa D, Garcia Dos Santos E and Giudicelli Y 2001. Site-specific control of rat preadipocyte adipose conversion by ovarian status: possible involvement of CCAAT/enhancer-binding protein transcription factors. *Endocrine* 15, 103–110.

Lipid accumulation in 3T3-L1 cells: effects of fatty acids and insulin

- Lee K, Hausman DB and Dean RG 1999. Expression of CCAAT/enhancer binding protein C/EBP α , β and δ in rat adipose stromal-vascular cells *in vitro*. *Biochimica et Biophysica Acta* 1450, 397–405.
- Li Y and Lazar MA 2002. Differential gene regulation by PPARgamma agonist and constitutively active PPARgamma2. *Molecular Endocrinology* 16, 1040–1048.
- Listenberger LL, Han X, Lewis SE, Cases S, Farese Jr RV, Ory DS and Schaffer JE 2003. Triglyceride accumulation protects against fatty acid-induced lipotoxicity. *Proceedings of the National Academy of Sciences of the United States of America* 100, 3077–3082.
- McNeel RL and Mersmann HJ 2003. Effects of isomers of conjugated linoleic acid on porcine adipocyte growth and differentiation. *Journal of Nutritional Biochemistry* 14, 266–274.
- McNeel RL, Smith EO and Mersmann HJ 2003. Isomers of conjugated linoleic acid modulate human preadipocyte differentiation. *In Vitro Cellular and Developmental Biology – Animal* 39, 375–382.
- Margetic S, Gazzola C, Pegg GG and Hill RA 2002. Leptin: a review of its peripheral actions and interactions. *International Journal of Obesity and Related Metabolic Disorders* 26, 1407–1433.
- Sakaue H, Ogawa W, Matsumoto M, Kuroda S, Takata M, Sugimoto T, Spiegelman BM and Kasuga M 1998. Posttranscriptional control of adipocyte differentiation through activation of phosphoinositide 3-kinase. *Journal of Biological Chemistry* 273, 28945–28952.
- Schneider N, Lanz S, Ramer R, Schaefer D and Goppelt-Strube M 2001. Up-regulation of cyclooxygenase-1 in neuroblastoma cell lines by retinoic acid and corticosteroids. *Journal of Neurochemistry* 77, 416–424.
- Shin SM, Kim K, Kim JK, Yoon SR, Choi I and Yang Y 2003. Dexamethasone reverses TGF-beta-mediated inhibition of primary rat preadipocyte differentiation. *Federation of European Biochemical Societies Letters* 543, 25–30.
- Spiegelman BM 1998. PPAR-gamma: adipogenic regulator and thiazolidinedione receptor. *Diabetes* 47, 507–514.
- Statistical Analysis Systems Institute 2002. SAS version 8.2. SAS Institute Inc., Cary, NC, USA.
- Thompson DL, Lum KD, Nygaard SC, Kuestner RE, Kelly KA, Gimble JM and Moore EE 1998. The derivation and characterization of stromal cell lines from the bone marrow of p53 $-/-$ mice: new insights into osteoblast and adipocyte differentiation. *Journal of Bone and Mineral Research* 13, 195–204.
- Tontonoz P, Hu E and Spiegelman BM 1994a. Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell* 79, 1147–1156.
- Tontonoz P, Hu E, Graves RA, Budavari AI and Spiegelman BM 1994b. mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes and Development* 8, 1224–1234.
- Torii SI, Kawada T, Matsuda K, Matsui T, Ishihara T and Yano H 1998. Thiazolidinedione induces the adipose differentiation of fibroblast-like cells resident within bovine skeletal muscle. *Cell Biology International* 22, 421–427.
- Wu Z, Bucher NL and Farmer SR 1996. Induction of peroxisome proliferator-activated receptor gamma during the conversion of 3T3 fibroblasts into adipocytes is mediated by C/EBPbeta, C/EBPdelta, and glucocorticoids. *Molecular and Cellular Biology* 16, 4128–4136.
- Wu Z, Xie Y, Morrison RF, Bucher NL and Farmer SR 1998. PPARgamma induces the insulin-dependent glucose transporter GLUT4 in the absence of C/EBPalpha during the conversion of 3T3 fibroblasts into adipocytes. *Journal of Clinical Investigation* 101, 22–32.
- Yamauchi T, Kamon J, Waki H, Murakami K, Motojima K, Komeda K, Ide T, Kubota N, Terauchi Y, Tobe K, Miki H, Tsuchida A, Akanuma Y, Nagai R, Kimura S and Kadowaki T 2001. The mechanisms by which both heterozygous peroxisome proliferator-activated receptor gamma (PPARgamma) deficiency and PPARgamma agonist improve insulin resistance. *Journal of Biological Chemistry* 276, 41245–41254.